

| | |
|--------------|---|
| Title | Comments on two vertebrate samples from early Islamic Jazirat al-Hulaylah (5th-9th c.AD) and Islamic Julfar (mid-14th-16th c. AD), United Arab Emirates |
| Author(s) | Beech, Mark |
| Citation | 金沢大学考古学紀要, 24: 197-203 |
| Issue Date | 1998-09-30 |
| Type | Departmental Bulletin Paper |
| Text version | |
| URL | http://hdl.handle.net/2297/1582 |
| Rights | |

*KURAに登録されているコンテンツの著作権は、執筆者、出版社（学協会）などが有します。

*KURAに登録されているコンテンツの利用については、著作権法に規定されている私的使用や引用などの範囲内で行ってください。

*著作権法に規定されている私的使用や引用などの範囲を超える利用を行う場合には、著作権者の許諾を得てください。ただし、著作権者から著作権等管理事業者（学術著作権協会、日本著作出版権管理システムなど）に権利委託されているコンテンツの利用手続については、各著作権等管理事業者に確認してください。

Comments on two vertebrate samples from early Islamic Jazirat al-Hulaylah (5th-9th c. AD) and Islamic Julfar (mid-14th - 16th c. AD), United Arab Emirates

Mark Beech

Department of Archaeology / Department of Biology (Environmental Archaeology Unit), University of York,

Introduction

This paper discusses two vertebrate samples retrieved from Early Islamic Jazirat al-Hulaylah and Islamic Julfar, Ras al-Khaimah emirate, United Arab Emirates. The sample of bones from Jazirat al-Hulaylah originate from context M3 (951125), which belongs to the earliest stratigraphic level at the site, level 1 (Sasaki 1995,1996; Sasaki and Sasaki 1996), and dates to the 5th-9th century AD. The sample from Julfar comes from 18 different ovens and a house context belonging to Level 6 (Sasaki 1993,1994), and dates to the mid-14th to 16th century AD. The purpose of this analysis was to provide an initial evaluation of the range of animals which were exploited at the two sites. Although the samples discussed here are small and only represent a small part of the total assemblage retrieved from the two sites, it seemed worthwhile making some preliminary comments on the basis of this evaluation. The author is shortly about to commence studying the remainder of the vertebrate assemblages from these two sites as part of his Phd research at the University of York, UK.

Methods

On-site dry sieving with 4mm mesh was used for all excavated contexts at both Jazirat al-Hulaylah and Julfar to permit the systematic recovery of animal bones. Both samples of bones were identified at the University of York using the comparative osteological collections of the Environmental Archaeology Unit as well as the author's personal osteological reference collection of Arabian Gulf marine fishes.

All fragments were counted and weighed (to the nearest gramme). Quantification was carried out as follows: In the case of crustacean remains only chelae survived and a simple count of the total number of fragments and weight was made (only counting chelae tips which were >50% complete). Fish remains were quantified using the method outlined by Beech (1997). In summary, the following elements were recorded: vomer, articular, dentary, maxilla, premaxilla, quadrate, hyomandibular, opercular, abdominal vertebra, caudal vertebra, vertebra (indeterminate), cleithrum, posttemporal, and otolith. In addition, a number of special elements were recorded which were distinctive of particular genera or species, e.g. scutes for Carangidae, and pharyngeals for Scaridae. The remains of marine turtle (Cheloniidae) were simply counted and weighed. This was because they mostly consisted of broken carapace fragments (with only the occasional rib or metapodial) and it was extremely difficult to define non-repeatable diagnostic parts of their skeleton. In the case of birds only the following skeletal parts were counted: proximal scapula,

proximal coracoid, distal humerus, proximal carpometacarpus, distal femur, distal tibiotarsus and distal tarsometatarsus. Terrestrial mammals were quantified using a modification of Davis (1992). The following parts of the skeleton were recorded: horncore (base with complete circumference intact), mandible (tooth row with at least 2 recordable teeth), isolated dP4s, P4s and molars, scapula (glenoid articulation), distal humerus, distal radius, carpal 2-3 (2 or 3 according to the taxon), distal metacarpus, ischial part of the acetabulum, distal femur, distal tibia, calcaneus, astragalus, distal metatarsus, proximal end of first phalanx, and third phalanx. At least 50% of a given part had to be present for it to be counted. Single metapodium condyles of caprines were counted as halves.

Counts were grouped into body zones in order to summarise the presentation of data. These were as follows: Crustacea: ch = chelum; Fishes: sk = neuro-/branchiocranium region: vomer, articular, dentary, maxilla, premaxilla, quadrate, opercular, pharyngeal., av = abdominal vertebra, cv = caudal vertebra, v = vertebra, indeterminate, ap = appendicular skeleton: cleithrum, posttemporal, oth = otolith, scute; Reptile: ca = carapace, rib = rib, mp = metapodial; Bird: wi = wing: scapula, coracoid, humerus, carpometacarpus, hl = leg: femur, tibiotarsus, tarsometatarsus; Mammal: sk = skull region: horncore, mandible, mandibular dP4, P4, M1/2 and M3, fl = forelimb: scapula, humerus, radius, carpal 2-3, metacarpus, hl = hindlimb: pelvis (acetabulum), femur, tibia, calcaneus, astragalus, metatarsus, ph = phalanges: first phalanx, third phalanx. All non-diagnostic bone fragments were grouped into the following categories: indeterminate fish, indeterminate bird and indeterminate mammal, and were counted and weighed. Information concerning the modern size range, habitat preferences and present day methods utilised to catch Arabian Gulf fishes are taken from Carpenter et al. (1997) and Randall (1995).

Results

Tables 1-2 present the results of this analysis.

Jazirat Al-Hulaylah

A total of 2164 bone fragments (3.38 kg) were recorded from Jazirat al-Hulaylah, out of which 1102 (51%) were identifiable to the level of family, genus or species.

At least two types of crabs were represented amongst the Crustacean remains: Portunidae (swimming crabs) and Xanthidae (stone crabs), of which the former were more numerous. In the Arabian Gulf there are three species within the Portunid family according to Carpenter et al. (1997), of which *Portunus pelagicus* (Linnaeus, 1758) is the largest in size and also the only species exploited commercially in the region at the present day. The xanthid crabs are generally much smaller than the portunids and can be found between the lower intertidal up to depths of 35m on rocky outcrops or coral reefs. They are only occasionally taken for human consumption at the present day. The majority of the crab fragments from Jazirat al-Hulaylah were burnt, presumably indicating their consumption.

Thirteen genera of fishes were represented. Important families which were exploited included the Carangidae (jacks), Sparidae (sea breams) and Scombridae (mackerel/tuna etc.). There was a significant concentration of sea bream fragments, which mostly consisted of dentaries and premaxillae belonging to the genus *Rhabdosargus*. These all belonged to fishes about 20-30cm in total length. Marine turtle was also eaten judging from the substantial number of burnt carapace fragments. Several of the *Phalacrocorax*

nigrogularis (Socotra cormorant) bones were also burnt indicating that these birds may also have been occasionally consumed. Domestic mammals were represented predominantly by the remains of caprines. Out of those bones which it was possible to distinguish between sheep (*Ovis ammon f. aries*) and goat (*Capra aegagrus f. hircus*), they all belonged to the latter. A single fragment was identified as belonging to camel. Interestingly this distal scapula (glenoid) fragment was markedly burnt thus perhaps also suggesting that they were sometimes eaten.

Julfar

A total of 1354 bone fragments (1.33 kg) were recorded from the various contexts at Julfar, out of which 438 (32%) were identifiable to the level of family, genus or species. No crustacea remains were identified within any of the Julfar contexts. Thirteen genera of fishes were represented. Important families which were exploited included the Carangidae (jacks) and Scombridae (mackerel/tuna etc.). Ovens 131 and 134 contained concentrations of fish scales presumably resulting from some form of processing activities. In the case of oven 131, the scales may belong to Haemulidae: *Pomadasys* sp. as within the same context there was an operculum with identical scales still attached to its outer surface. Marine turtle only occurred as occasional carapace fragments in two out of the 19 sampled contexts. Both sheep and goat were represented amongst the caprine remains. A single fragment of a dog pelvis occurred in oven 134. This did not exhibit any signs of butchery or burning suggesting that it may have been eaten.

Discussion

The diverse range of fishes exploited at both sites suggests that subsistence strategies were geared towards exploiting a multi-species based fishery. Fish bones make up the bulk of the material at both Jazirat al-Hulaylah and Julfar perhaps suggesting their relative dietary importance. Caprines, mainly goats, would have provided regular contributions of meat as well as milk and other secondary products. Camels would have been used as a mode of transport and as beasts of burden for the shipping of materials, although it seems they may have been sometimes eaten during early Islamic Jazirat Al-Hulaylah. The vertebrate assemblages from Jazirat al-Hulaylah and Julfar broadly match those from the nearby site of Kush in terms of the importance of ovicaprid husbandry as well as fishing (Beech and Pipe 1997). Although the samples are small, the fact that a range of ovicaprid body parts are represented suggests that whole animals were brought to the site "on the hoof" for slaughter and consumption within the settlement area.

Table 3 summarises the modern size range, habitat preferences and present day methods utilised to catch the fish groups represented at the two sites. This suggests that fishing was predominantly carried out at both sites in adjacent coastal inshore waters near to coral reefs. Most of the fish represented were probably caught using a combination of inshore nets, traps and handlines. Some fishing however was also done probably further offshore in boats to target some of the pelagic species like tuna and kingfish (*Thunnus* sp. and *Scomberomorus* sp.).

Further more detailed work on the remainder of the vertebrate assemblages from Jazirat al-Hulaylah and Julfar will undoubtedly highlight further details concerning the economic basis of the early and later Islamic periods of the region.

References

- Beech, M. (1997). *Ancient Marine Resource Exploitation in the southern Arabian Gulf: An Archaeozoological Perspective*. Unpublished Dphil. TAP report, submitted 31st October, Departments of Archaeology and Biology, University of York.
- Beech, M. and Pipe, A. (1997) The Animal Bones. In: Kennet, D., Kush: a Sasanian and Islamic-period archaeological tell in Ras al-Khaimah (U.A.E.). *Arabian Archaeology and Epigraphy* 8: 284-302 (pp.297-298).
- Carpenter, K. E., F. Krupp, D.A. Jones, and U. Zajonz (1997). *FAO Species Identification Field Guide for Fishery Purposes: The Living Marine Resources of Kuwait, Eastern Saudi Arabia, Bahrain, Qatar, and the United Arab Emirates*. FAO: Rome.
- Davis, S. (1992). *A rapid method for recording information about mammal bones from archaeological sites*. London, AML report 19/92.
- Randall, J.E. (1995). *Coastal Fishes of Oman*. Crawford House: Bathurst, Australia.
- Sasaki, T. (1993). "Excavations at Julfar in 1992 season." *Bulletin of Archaeology - The University of Kanazawa* 20: 1-49.
- Sasaki, T. (1994). "1993 excavations at Julfar." *Bulletin of Archaeology - The University of Kanazawa* 21: 1-106.
- Sasaki, T. (1995). "1994 excavations at Jazirat al-Hulayla, Ras al-Khaimah." *Bulletin of Archaeology - The University of Kanazawa* 22: 1-74.
- Sasaki, T. (1996). "Umayyad and Abbasid finds from the 1994 excavations at Jazirat al-Hulayla." *Bulletin of Archaeology - The University of Kanazawa* 23: 179-222.
- Sasaki, T. and H. Sasaki (1996). "1995 excavations at Jazirat al-Hulayla, Ras al-Khaimah." *Bulletin of Archaeology - The University of Kanazawa* 23: 37-178.

| FAMILY | TAXON | JHU | JU | TYPICAL LENGTH (CM) | HABITAT | MODERN FISHERIES CAPTURE METHOD |
|-----------------|---------------------------|-----|----|---------------------|--|--|
| Carcharhinidae | Carcharhinus sp. | * | * | 100-340 | Inshore - offshore | Gillnet, line gear, longline |
| Myliobatidae | cf. Aetobatus narinari | * | | up to 230cm (width) | Open water | Drift net, Gillnet |
| Clupeidae | | | * | 10-25 | Coastal pelagic | Seine, gillnet, set net, lift net, shallow trawl |
| Belonidae | | * | | 30-124 | Pelagic offshore- pelagic coastal | Surface casting/trolling, seine, drift net, |
| Platycephalidae | Platycephalus indicus | * | | 60-100 | Benthic on sand or mud bottoms, shallow to 25m | Bottom trawl |
| Serranidae | Epinephelus sp. | * | * | 40-157 | Seagrass beds, coarse sand, rocky banks, coral reefs, 6-200m | Hook-and-Line, trap, trawl |
| Carangidae | Carangoides sp. | * | * | 50-100 | Coastal inshore, rocky bottom, coral reef | Hook-and-Line, trap, gillnet, spear, trawl |
| Carangidae | Caranx sp. | | * | 60-165 | Coastal waters, rocky reefs, coral reefs | Hook-and-Line, trap, gillnet, spear, purse seine |
| Carangidae | cf. Gnathanodon speciosus | * | | Up to 110 | Inshore rocky reefs, deep lagoons, seaward reefs | Gillnet, spear |
| Gerreidae | Gerres sp. | | * | 15-35 | Clear, shallow waters, sandy bottom to 50m | Bottom trawl, beach seine |
| Lutjanidae | Lutjanus sp. | * | | 20-100 | Estuarine, coral reefs, rocky reefs, at 3-100m | Handline, trap, bottom longline, trawl, gillnet |
| Haemulidae | Pomadasys sp. | ?* | * | 25-80 | Coastal, sand and mud bottom, rocky or rock-sand bottom, to 60m | Trawl, trap, handline, gillnet |
| Nemipteridae | Scolopsis sp. | | * | 13-28 | Shallow sandy or mud bottom, near coral reefs, to 60m | Trap, gillnet, trawl |
| Lethrinidae | Lethrinus sp. | * | * | 20-80 | Coastal over sand and hard bottoms, on/near reefs and seagrass beds, 1-80m | Trap, handline, trawl |
| Sparidae | Acanthopagrus sp. | * | | 30-75 | Coastal areas, rough and muddy sand bottom, coral reefs, shallow to 50m | Bottom trawl, trap, handline |
| Sparidae | Rhabdosargus sp. | * | * | 35-60 | Coral reef, sandy or mud-sandy bottoms, shallow to 60m | Gillnet, seine, trap, handline, trawl |
| Sparidae | Argyrops spinifer | | * | 30-65 | Variety of bottoms, 5-100m | Trawl, trap |
| Scaridae | Scarus sp. | * | | 32-57 | Coral reefs, lagoons, shallow water over sand or sea grass bottoms, 1-60m | Trap, small-scale net gear |
| Sphyraenidae | Sphyraena sp. | * | * | 20-180 | Near surface, close inshore over shallow banks, near bottom, to 100m | Trolling line, nets |
| Scombridae | Scomberomorus sp. | | * | 55-220 | Epipelagic coastal/neritic | Driftnet, trawl and trolling |
| Scombridae | Thunnus sp. | * | * | 70-200 | Epipelagic oceanic | Longline, purse seine |

TABLE 3.

Fishes represented at Jazirat Al-Hulayla (JHU) and Julfar (JU) and their typical modern size ranges, preferred habitats and modern fisheries methods utilised to catch them - data taken from Carpenter et al. (1997) and Randall (1995).

* = present.