


# TRIBULUS



Bulletin of the Emirates Natural History Group

Vol. 9.1

SPRING 1999



## NOTES FOR CONTRIBUTORS

TRIBULUS is the name of the Bulletin of the Emirates Natural History Group. The Group was founded in 1976, and over the next fourteen years, 42 issues of a duplicated Bulletin were published. The revised format of TRIBULUS, introduced in 1991, permits the inclusion of black and white and colour photographs, not previously possible.

TRIBULUS is published twice a year, in April and October. The aim of the publication is to create and maintain in standard form a collection of recordings, articles and analysis on topics of regional archaeology and natural history, with the emphasis on the United Arab Emirates and adjacent areas. Papers, short notes and other contributions are welcomed from anyone but should not have been published elsewhere. Guidelines are set out below. The information carried is as accurate as the Editorial Board and Advisory Panel can determine, but opinions expressed are those of the authors alone.

Correspondence and enquires should be sent to:


The Editor,  
TRIBULUS,  
Emirates Natural History Group,  
P.O. Box 2380,  
Abu Dhabi - U.A.E.

### Editorial Board:

*H.E. Sheikh Nahayan bin Mubarak Al Nahayan, Patron,*  
*Rob Western, Chief Editor,*  
*Peter Hellyer, Managing Editor,*  
*Simon Aspinall, Deputy Editor,*  
*Dr. Michael Gillett*  
*Steve James*

### Advisory Panel:

*Professor Jens Eriksen (Life Sciences)*  
*Professor Graham Evans (Earth Science)*  
*Professor Dan Potts (Archaeology)*  
*Dr. Saif Al Ghais (Marine Science and Conservation)*  
*Dr. David Jones (Fauna)*  
*Dr. Geoffrey R.D. King (Archaeology)*



The plant motif above is of the genus *Tribulus*, of which there are six species in the UAE. They all have pinnate leaves, yellow flowers with free petals and distinctive five-segmented fruits. They are found throughout the country, except in coastal *sabkha*.

The animal motif above is of a tiny golden bull, excavated from the early Second Millennium grave at Qattarah, Al Ain. The original is on display in Al Ain Museum, and measures above 5 cm by 4 cm.

Manuscripts should be typed, on one side only, and double-spaced, and should be accompanied by a disc for material in excess of 500 words in length. A short abstract should precede the article, with the address(es) of the author(s) at the end.

Photographs may be submitted and should be either glossy black-and-white or colour prints or colour slides, which should be clearly captioned. Line drawings and maps should be in black ink on strong white or translucent paper.

References should give the author's name, with the year of publication in brackets, and with the list of articles, showing title and publisher, in date order.

Scientific names should follow customary nomenclature in Latin, while the English and, if appropriate, available Arabic names should also be supplied.

**International Standard Serial Number (ISSN): 1019 - 6919.**

# TRIBULUS

Vol. 9.1 Summer 1999

## Contents

	Page
Editorial .....	4
Freshwater Snails of the UAE.....	5
by Gary R. Feulner and Stephen A. Green	
An AMS radiocarbon chronology for the late Umm an-Nar type tomb at Tell Abraq .....	9
by D.T. Potts and Lloyd Weeks	
Sperm collection and oocyte retrieval in the Arabian Leopard <i>Panthera pardus nimr</i> .....	11
by Peter Wright	
The Finless Porpoise <i>Neophocaena phocaenoides</i> in the Arabian Gulf .....	13
by Simon Aspinall and Robert Baldwin	
Clam Shrimps (Branchiopoda, Crustacea) - a new discovery for the UAE .....	16
by Richard Hornby	
Notes and Queries .....	18
Arabia's oldest date stones; a new species of trigger fish for the Arabian Gulf; Conference notes; Research protocol for Endangered Arabian Wildlife Centre; Records of owl flies in the Al Ain area; Notes on newly recorded butterflies from UAE and northern Oman; A first record of Blackstart for the UAE; A first confirmed record of the Leopard Butterfly from Sharjah Emirate; Memorial unveiled to UAE's first oil well.	
Recent publications/Reviews .....	27
Editorial Notebook .....	29
Archaeological Review .....	30
Recorders' Reports: Mammals and Miscellaneous .....	31
Bird Report .....	32
Cover illustrations:	
Front: undersurface, UNS, of a small male <b>Pontia glauconome</b> - dry season form (top); upper surface, UPS, and UNS of male and female <b>P. glauconome</b> - wet season form (row 2); UNS and UPS of male and female <b>P. glauconome</b> - extreme dry season form (row 3); UPS of male <b>Pontia daplidice</b> extreme dry season form. (see article on P. 22).....	Picture by Ashok Prasad
Back: A <b>Hottentota jayakari</b> with two tails ,from the Wadi Qawr.....	Picture by Dr. J. Kinne

The Editorial Board of TRIBULUS and the Committee of the Emirates Natural History Group acknowledge, with thanks, the support of the Group's Corporate members, a full list of whom can be found on Page 4, without whom publication in this format would be impossible. We also acknowledge the support and encouragement of our Patron, H.E. Sheikh Nahayan bin Mubarak Al Nahayan, the U.A.E. Minister of Higher Education and Scientific Research.

TRIBULUS is published for circulation to members of the ENHG, and is also available for subscribers in the UAE and overseas or through the Ornithological Society of the Middle East. Details on request.

The text of this issue is also available on the website of the UAE Ministry of Information and Culture: [www.uaeinteract.com](http://www.uaeinteract.com)

Published by the Emirates Natural History Group, P.O. Box 2380, Abu Dhabi, United Arab Emirates.

The Group is a non-governmental member of IUCN, the World Conservation Union.

Printed for the ENHG by Dar Al Fajr Press, Printing, Publishing and Advertising,

P.O.Box 505, Abu Dhabi, UAE



## NOTES AND QUERIES

### ***Dalma archaeological site yields Arabia's oldest date stones***

The results of radiocarbon dating of two date *Phoenix dactylifera* stones found by the Abu Dhabi Islands Archaeological Survey, ADIAS, on Dalma has shown that they are the oldest ever found in Arabia. The results suggested that one date stone can be dated to 4670 (+/-130) BC, and the other to 5110 (+/-160) BC, that is about 6,500 - 7,000 years ago.

The dating was carried out at the University of Arizona working in collaboration with the Scottish Universities Research and Reactor Centre radiocarbon laboratory at the University of Glasgow, in Scotland.

Both date stones, together with impressions of date stones on fragments of mudbrick, were recovered during excavations on a site in the compound of the Abu Dhabi Women's Association branch on Dalma, where ADIAS has been working for several years. They were identified during the course of work organised in early 1998 with the support of Minister of Information and Culture Sheikh Abdulla bin Zayed Al Nahyan.

The two Dalma date stones represent the oldest radiometrically dated evidence yet available for the consumption of dates within the Gulf region, as well as probably some of the earliest evidence of the date palm found anywhere in the whole of the Middle East. Previously, the earliest evidence for date palm remains in the UAE was excavated from the Hili 8 site in Al Ain, dated to around 3,000 BC. The Dalma date stones are at least 1,500 years, and perhaps over 2,000 years older.

The two date stones also represent some of the earliest remains of date consumption found within the entire Middle East. Although it cannot be determined if they represent wild or cultivated dates, they certainly confirm that dates were being consumed at this early time.

Other finds from the Dalma site include at least two round house-like structures with surviving post-holes and floors, one of which is at least 7 metres in diameter. There are also small quantities of imported painted pottery from the 'Ubaid culture of southern Mesopotamia. Since Dalma was already almost certainly an island at the time, the 'Ubaid pottery must have made at least part of its journey by sea, offering us the first confirmed evidence of the maritime trading connections of the people of the Emirates.

Large quantities of what appear to be locally made gypsum plaster vessels of a type not known anywhere else in the Middle East have also been found. During an inspection visit to the Dalma site at the end of March, a large portion of one of these vessels was found which is the most complete vessel of this period ever identified in the Emirates.

The Dalma site has also yielded thousands of Late Stone Age flint flakes and a number of stone tools; other finds included ornamental beads, and huge quantities of food debris in the form of marine shells and animal and fish bones.

**Mark Beech**

### ***A new species of trigger fish recorded for the Arabian Gulf***

**Teleostei, Tetraodontiformes, Balistidae, *Canthidermis maculatus* (Bloch, 1786) – the Spotted Oceanic Triggerfish**

This note describes an unusual triggerfish purchased by the author in Ras Al Khaimah fish *souq* on 12 April 1998 which has been identified as the **Spotted Oceanic Triggerfish** *Canthidermis maculatus*. The record is the first for the Arabian Gulf. The fish had 3 dorsal hard spines and 23 soft dorsal rays, as well as 21 soft anal rays. Measurements were taken as follows:

**Total length (TL):** measured from the tip of the snout to the end of the upper lobe of the caudal fin = 33.5 cm.

**Standard length (SL):** measured from the tip of the snout to the end of the axial skeleton (determined by bending the tail upwards) = 28 cm.

**Head length (HL):** from the tip of the snout to the posterior edge of the gill cover (operculum) = 9 cm.

**Body depth (BD):** the deepest point from the origin of the dorsal fin vertically downwards to the pelvic fin = 11 cm.

The fresh capture weight of the fish was 735 g. Its colour was dark purplish to black with white patches on the lower ventral sides of its body. The caudal fin was slightly rounded with small notches present in moderately pronounced lobes.

Three triggerfish species commonly occur in the Arabian Gulf according to the most recently published surveys (Carpenter *et al.*, 1997; Randall, 1995). These are:

- the **starry triggerfish** *Abalistes stellatus* (Lacepède, 1798),
- the **picasso triggerfish** *Rhinecanthus assasi* (Forsskal, 1775),
- the **flagtail triggerfish** *Sufflamen chrysopterus* (Bloch and Schneider, 1801).

The specimen from Ras Al Khaimah does not however match with any of these. Its distinctive colour as well as spine/ray counts and body/fin shapes preclude it from belonging to any of these previously recorded species. Outside the Gulf, in adjacent Omani waters, a number of other triggerfishes are known to also occur (Randall, 1995). These include:

- the **largescale triggerfish** *Canthidermis macrolepis* (Boulenger, 1887),
- the **Indian Ocean durgon** *Melichthys indicus* (Randall and Klauswitz, 1973),
- the **redtooth triggerfish** *Odonus niger* (Rüppell, 1836),
- the **bridled triggerfish** *Sufflamen fraenatus* (Latreille, 1804).

These latter three species have quite separate distinct colour and body/fin shapes. The Ras Al Khaimah triggerfish looks closest to the **largescale triggerfish** *Canthidermis macrolepis*, although there are a number



of clear differences. *Canthidermis macrolepis* has quite an elongate sort of body for a triggerfish, and is grey in colour, shading to pale grey ventrally. The edges of its second dorsal, anal and caudal fins are blackish. Its caudal fin is double emarginate with produced lobes (Randall, 1995: 393; Gill and Randall, 1997).

A closer match for the specimen purchased in Ras Al Khaimah souq is another species within the same genus as the largescale triggerfish, namely, the **spotted oceanic triggerfish** *Canthidermis maculatus* (Bloch, 1786), with which the specimen can be confidently identified.

This is a triggerfish which occurs circumglobally in both tropical and temperate seas. It has been recorded previously within the Western Indian Ocean (Smith and Heemstra, 1986). The head, body and fins of adults are dark; their body has elongated white spots that may disappear with growth (Smith and Heemstra, 1986). This species is epipelagic through nearly all its life, and is often associated with drifting objects (Myers, 1991). It can occur on deep rocky slopes (Robins and Ray, 1986), and appears to school in large numbers based on catches from set nets (Masuda *et al.*, 1975). In some parts of the world it is marketed fresh (Matsuura, 1997). It commonly occurs at a length of around 30 cm (Bussing, 1995), but can occur up to a maximum length of 50 cm (Edwards, 1990).

According to FishBase98 (FishBase, 1998; Froese and Pauly, 1998), three of the localities nearest to the Arabian Gulf with records of *Canthidermis maculatus* are:

- Obock, Djibouti, in the Gulf of Aden, near the entrance to the Red Sea. Location: 11°58' N 043°20' E (a 56cm TL dry mounted specimen collected by Maindron, retained in the Musée National d'Histoire Naturelle, Paris - MNHN 18930072).

- Maldivé Islands, Indian Ocean. Location: 03°40' N 073°49' E (recorded in the fish list for the Maldives by: Randall and Anderson, 1993).


- B.esperance(c.), South Africa. Location: 34°21' S 18°25' E (a 30.8 cm TL dry mounted specimen collected by Verreaux, retained in the Musée National d'Histoire Naturelle, Paris - MNHN A8511).

The discovery of **spotted oceanic triggerfish** within the Ras Al Khaimah fish market suggests that the northern part of its range in the Western Indian Ocean may be more extensive than previously realised. Fishing boats venturing out from Ras Al Khaimah rarely stay away from port for more than two days at a time, and the fishing grounds exploited are almost completely Arabian Gulf-based. Sailing around the Musandam peninsula involves too much time and is not deemed to be economically viable. It seems likely, therefore, that the specimen may have originated from the waters adjacent to Ras Al Khaimah, although it is not possible to be certain whether it was caught in the territorial waters of the UAE or of Iran or Oman, which also share control of waters around the Straits of Hormuz. Juvenile triggerfishes are, however, often found in association with floating debris and vegetation, so they may have drifted into adjacent regions some distance from their native territory.

#### Acknowledgements:

My 1998 field research trip to the UAE was financed by The British Council (Abu Dhabi), ERWDA (Environment Research and Wildlife Development Agency, Abu Dhabi), ADIAS (Abu Dhabi Islands Archaeological Survey) and the University of York. Accommodation and subsistence in Ras Al Khaimah was kindly provided by Sheikh Sultan bin Saqr al-Qassimi, Director of the Department of Antiquities and Museums of Ras al-Khaimah. Christian Velde, resident archaeologist at the National Museum of Ras al-Khaimah, also provided valuable support to the author during his stay.

Callum Roberts and Julie Hawkins (Environment Department, University of York) generously provided access to FishBase98 and were helpful in providing



Spotted Oceanic triggerfish from Ras Al Khaimah souq. Picture by Mark Beech.



other bibliographic references. Both Tony Gill (Natural History Museum, London) and Jack Randall (Bishop Museum, Hawaii) gave advice concerning the identification of the triggerfish. Tony Gill kindly provided me with a copy of his and Randall's recent paper on *Canthidermis macrolepis*.

## References

Bussing, W.A. (1995). Balistidae. p. 905-909. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and V. Niem (eds.) Guia FAO para Identificación de Especies para lo Fines de la Pesca. Pacifico Centro-Oriental. 3 Vols. FAO, Rome.

Carpenter, K. E., Krupp, F., Jones, D.A. and Zajonz, U. (1997). FAO species identification guide for fishery purposes. The Living Marine Resources of Kuwait, Eastern Saudi Arabia, Bahrain, Qatar, and the United Arab Emirates. Rome, FAO.

Edwards, A. (1990). Fish and fisheries of Saint Helena Island. Centre for Tropical Coastal Management Studies, University of Newcastle upon Tyne, England.

FishBase. (1998). FishBase 98 CD-ROM. ICLARM, Manila.

Froese, R. and D. Pauly (eds.) (1998). FishBase 98: concepts, design and data sources. ICLARM, Manila, Philippines.

Gill, A.C. and Randall, J.E. (1997). Redescription of and lectotype designation for *Balistes macrolepis* Boulenger, 1887, a senior synonym of *Canthidermis longirostris* Tortonese, 1954 and *C. villosus* Fedoryako, 1979 (Teleostei, Tetraodontiformes, Balistidae). Bull. Nat. Hist.

Mus. Lond. (Zool.) 63(1): 27-31.

Masuda, H., C. Araga and T. Yoshino. (1975). Coastal fishes of Southern Japan. Tokai University Press, Tokyo, Japan.

Matsuura, K. (1997). Balistidae. Triggerfishes. In K.E. Carpenter and V. Niem (eds.) FAO Identification Guide for Fishery Purposes. The Western Central Pacific. (in prep).

Myers, R.F. (1991). Micronesian reef fishes. Second Ed. Coral Graphics, Barrigada, Guam.

Randall, J. E. (1995). Coastal Fishes of Oman. Bathurst, Australia, Crawford House Publishing.

Randall, J.E. and C. Anderson (1993). Annotated checklist of the epipelagic and shore fishes of the Maldives Islands., Ichthyol. Bull. of the J.L.B. Smith Inst. of Ichthyol. 59:47.

Robins, C.R. and G.C. Ray. (1986). A field guide to Atlantic coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A.

Smith, M.M. and P.C. Heemstra. (1986). Balistidae. p. 876-882. In M.M. Smith and P.C. Heemstra (eds.) Smiths' Sea Fishes. Springer-Verlag, Berlin.

Mark Beech  
Department of Archaeology  
The King's Manor  
University of York  
York YO1 2EP, UK  
E-mail: mjb117@york.ac.uk

---

## CONFERENCE NOTES

---

### Reclaiming the Past - Shaping the Future

A one day symposium with the above title was hosted by the American University of Sharjah on February 11th 1999. Sub-titled "A symposium on material culture in the UAE," it was organised by the AUS School of Architecture and Design, and focussed primarily on recent research into the country's archaeology and on traditional architecture and its restoration.

With participants including many of the familiar names in Emirates archaeology, the Symposium provided a useful opportunity for discussion on ways in which university students could become involved in the study of the country's heritage. AUS is now planning to launch a special course in 'heritage management.' Proceedings are scheduled to be published later this year.

### 1st Abu Dhabi International Arabian Oryx Conference

Hosted by the Environment & Wildlife Management Unit of the Private Department of H.H. Sheikh Zayed bin Sultan Al Nahyan, the conference was held in Abu Dhabi in February 1999, and concluded with the drafting of a memorandum of understanding, the 'Abu Dhabi Declaration,' essentially an agreement between interested parties to foster increased collaboration between breeding establishments and research institutes worldwide and between environmental authorities in the range states. The facilitation of

successful reintroductions to the wild was an agreed major goal of attending parties. A regional advisory group is to be set up, with Oman having offered to host the first meeting of such a group. Proceedings will be published later this year. Contact: The Director, EWM, P.O. Box 77, Abu Dhabi, UAE. Fax: 00-971 +2 - 663033. Email: wildlife@dpanet.org.ae.

### ECO-ARABIA '99

Sponsored by Shell, a two day 'Eco-Arabia '99' conference in Dubai in April examined protection of the marine and coastal environments of the Middle East. Speakers, the majority from overseas but many from within the GCC, were from governmental ministries and research institutes, private business and public sector industry. Partnership, collaboration and communication were identified as key issues of mutual benefit.

### 'Arab Envirotec-1' Conference

The First Arab International Conference and Exhibition on Environmental Biotechnology, organised by the Emirates Heritage Club and under the patronage of UAE Deputy Prime Minister Sheikh Sultan Bin Zayed Al Nahyan, was held in Abu Dhabi from 5-8 March 1999. A series of technical and non-technical presentations were delivered.

A concurrent exhibition featured displays by local commercial companies and non-profit-making environmental NGOs.